



UNIVERSITI PUTRA MALAYSIA

**STUDIES ON THE IMPROVEMENT OF THE NUTRITIVE VALUE AND
UTILIZATION OF PALM KERNEL CAKE AS A FEED
RESOURCE FOR RUMINANTS**

CHARURAT CHINAJARIYAWONG

IB 2000 3

**STUDIES ON THE IMPROVEMENT OF THE NUTRITIVE VALUE AND
UTILIZATION OF PALM KERNEL CAKE AS A FEED
RESOURCE FOR RUMINANTS**

By

CHARURAT CHINAJARIYAWONG

**Thesis Submitted in Fulfillment of the Requirements for the Degree of
Doctor of Philosophy in the Institute of Bioscience
Universiti Putra Malaysia**

March 2000



Abstract of thesis submitted to the Senate of Universiti Putra Malaysia
in fulfilment of the requirements for the Degree of Doctor of Philosophy

**STUDIES ON THE IMPROVEMENT OF THE NUTRITIVE VALUE AND
UTILIZATION OF PALM KERNEL CAKE AS A FEED
RESOURCE FOR RUMINANTS**

By

CHARURAT CHINAJARIYAWONG

MARCH 2000

Chairman: Professor Tan Sri Dato' Syed Jalaludin, Ph.D.

Faculty : Institute of Bioscience

Utilization of palm kernel cake (PKC) as the principal feed for ruminants was investigated in four studies. The study on the nutritive value showed that PKC-solvent extracted (PKC-S) and PKC-expeller pressed (PKC-E) contained similar amounts of OM, CP, CF and ADL. PKC-E had higher DM, EE and GE but lower NFE, NDF, ADF and shell contents. Amounts of Ca, P, Cu and Zn were similar in both types of PKC, but Na was higher and Mg was lower in PKC-S. Ruminal pH, NH₃-N concentration, molar proportions of VFA and passage rates of liquid and small digesta particles in cattle fed either PKC-S or PKC-E were similar. Total VFA concentration was higher in cattle fed PKC-S. Effective DM and OM degradabilities were lower in animals fed PKC-S, but degradation

rates were similar. PKC-E contained more soluble fractions than PKC-S. CP of PKC-S was more soluble, but its effective degradability and degradation rate were similar to that of PKC-E.

The study on the effects of formaldehyde treatments on CP degradability of both types of PKC showed that CP degradability decreased with increasing levels of formaldehyde. Acid-pepsin digestibility showed no difference in CP digestibility among untreated and formaldehyde-treated PKC up to 2.5g/100g CP.

The study on the effects of PKC and formaldehyde treatment on the performance and nutrient utilization of cattle showed that feed intake was lower in cattle fed PKC-E, but growth rates and feed per gain ratios did not differ from those fed PKC-S. Formaldehyde-treated (2g/100g CP) PKC gave poorer growth rates and feed conversion ratios when compared to untreated PKC. Types of PKC and formaldehyde treatment did not affect the nutrient digestibility, but PKC-E gave lower N retention. Formaldehyde had no effect on N retention.

In the final study, lambs fed PKC + 15% grass showed improvement in intakes and growth rates. Urea at either 0.5 or 1.0%, had no effect on feed intakes and feed conversion ratios. However, growth rates of lambs fed PKC + 0.5% urea were higher compared to other groups. Lambs fed PKC plus 15% grass and 0.5% urea had the highest intake and growth rate. Carcass traits were not affected by urea but grass supplementation reduced dressing percent and kidney fat of lambs.

Abastrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Doktor Falsafah

**KAJIAN KE ATAS PENINGKATAN PEMAKANAN DAN PENGGUNAAN
MEAL ISIRONG KELAPA SAWIT SEBAGAI SUMBER MAKANAN UNTUK
RUMINAN**

Oleh

CHARURAT CHINAJARIYAWONG

MAC 2000

Pengerusi: Profesor Tan Sri Dato' Syed Jalaludin Syed Salim, PhD

Fakulti : Institut Biosains

Kajian penggunaan meal isirong kelapa sawit (PKC) sebagai komponen utama ruminan telah dijalankan. Dari kajian nilai pemakanan menunjukkan nilai pemakanan 'PKC-solvent extracted' (PKC-S) dan 'PKC-expeller extracted' (PKC-E) mengandungi kandungan OM, CP, CF and ADL yang sama. PKC-E mengandungi jumlah DM, EE dan GE yang tinggi tetapi NFE, NDF, ADF dan kandungan tempurung yang rendah. Jumlah Ca, P, Cu dan Zn pada kedua-dua jenis PKC adalah sama tetapi amaun Na lebih tinggi dan Mg lebih rendah pada PKC-S. Nilai pH rumen, kepekatan $\text{NH}_3\text{-N}$, peratus molar VFA, kadar pengaliran bendalir dan partikel halus pada lembu yang diberi makan PKC-S dan PKC-E adalah sama. Jumlah kepekatan VFA lebih tinggi pada lembu yang diberi PKC-S. Degradasi efektif DM dan OM lebih rendah pada haiwan yang diberi PKC-S tetapi kadar degradasi sama. PKC-E mempunyai nilai fraksi larut yang lebih tinggi dari PKC-S. CP PKC-S adalah lebih larut daripada PKC-E tetapi degradasi efektif dan kadar degradasi kedua-dua PKC adalah sama.

Kajian kesan formaldehid ke atas kadar degradasi CP bagi PKC-S dan PKC-E menunjukkan degradasi CP menurun dengan peningkatan tahap formaldehid. Penghadaman asid-pepsin menunjukkan tiada perbezaan pada penghadaman CP di antara PKC yang tidak dirawat dan PKC yang dirawat dengan formaldehid hingga 2.5g/100g CP.

Dalam kajian kesan PKC dan rawatan formaldehid ke atas prestasi dan pengambilan makanan pada lembu menunjukkan kadar pengambilan makanan yang rendah pada lembu yang diberi PKC-E tetapi kadar pertumbuhan dan nisbah makanan pertumbuhan tidak berbeza daripada lembu yang diberi PKC-S. Lembu yang diberi makan PKC yang dirawat formaldehid pada tahap 2g/100g CP menunjukkan kadar pertumbuhan dan nisbah penukaran makanan yang merosot dibandingkan dengan lembu yang diberi PKC tanpa rawatan. Jenis PKC dan rawatan formaldehid tidak memberi kesan ke atas penghadaman nutrien tetapi lembu yang diberi PKC-E menunjukkan retensi nitrogen yang rendah. Rawatan formaldehid tidak memberi kesan terhadap retensi nitrogen.

Kajian terakhir menunjukkan peningkatan pengambilan makanan dan kadar pertumbuhan anak bebiri yang diberi PKC dan 15% rumput. Urea pada tahap 0.5 atau 1.0% tidak memberi kesan terhadap pengambilan makanan dan nisbah penukaran makanan. Walaubagaimana pun, kadar pertumbuhan anak bebiri yang diberi PKC dan 0.5% urea lebih tinggi dibandingkan dengan dengan kumpulan yang lain. Anak bebiri yang diberi makan PKC serta 15% rumput dan 0.5% urea menunjukkan pengambilan makanan dan pertumbuhan yang tinggi. Ciri karkas tidak terjejas dengan penambahan urea tetapi penambahan rumput mengurangkan peratusan berat bersih dan lemak ginjal anak bebiri.

ACKNOWLEDGMENTS

I would like to express my appreciation and sincere gratitude to my supervisors Professor Tan Sri Dato' Dr. Syed Jalaludin bin Syed Salim, Professor Dr. Ho Yin Wan and Associate Professor Dr. Norhani bt. Abdullah of the Institute of Bioscience, Universiti Putra Malaysia for their invaluable guidance and advice throughout the course of this study and in the preparation of this thesis.

I would like to thank the Prince of Songkla University for granting me the study leave. Sincere appreciation is also expressed to the Southeast Asian Regional Centre for Graduate Study and Research in Agriculture (SEARCA) for providing me a scholarship, and the Department of Animal Sciences, Faculty of Veterinary Medicine and Animal Sciences, for providing the research facilities.

I am also much indebted to Professor Dr. Rudy I. Hutagalung, my former supervisor, for his invaluable guidance and advice at the beginning of my study.

I am also most grateful to Professor Dato' Dr. Mohd. Mahyuddin bin Mohd. Dahan for his invaluable guidance in carcass measurement.

I would like to thank Associate Professor Dr. Zainal Aznam bin Mohd. Jelani for preparing rumen fistulated animals for use in this study.

Appreciation is extended to all the staff of Animal Science Department for their corporation and assistance.

Finally, I wish to express my deepest gratitude to my family for their encouragement.

I certify that an Examination Committee met on 9 March, 2000 to conduct the final examination of Charurat Chinajariyawong on her Doctor of Philosophy thesis entitled " Studies on the Improvement of the Nutritive Value and Utilization of Palm Kernel Cake as a Feed Resource for Ruminants" in accordance with Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Prof. Madya Dr. Abdul Razak Alimon
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Prof. Tan Sri Dr. Syed Jalaludin Syed Salim
Vice Chancellor
Universiti Putra Malaysia
(Member)

Prof. Dr. Ho Yin Wan
Institute of Bioscience
Universiti Putra Malaysia
(Member)


Prof. Madya Dr. Norhani Abdullah
Faculty of Science and Environmental Studies
Universiti Putra Malaysia
(Member)



MOHD. GHAZALI MOHAYIDIN, Ph.D
Professor/Deputy Dean of Graduate School

Date: **10 MAR 2000**

This thesis was submitted to the Senate of Universiti Putra Malaysia and was accepted as fulfillment for the degree of Doctor of Philosophy.


KAMIS AWANG, Ph.D.
Associate Professor,
Dean of Graduate School,
Universiti Putra Malaysia
Date: 11 MAY 2000

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or currently submitted for any other degree at UPM or other institutions.

Charurat Chinarajariyawong

(CHARURAT CHINAJARIYAWONG)

Date: 9 / 3 / 2000

TABLE OF CONTENTS

	Page
ABSTRACT	ii
ABSTRAK	iv
ACKNOWLEDGEMENTS.....	vi
APPROVAL SHEETS.....	viii
DECLARATION FORM.....	x
TABLE OF CONTENTS.....	xi
LIST OF TABLES.....	xv
LIST OF FIGURES.....	xx
LIST OF ABBREVIATIONS.....	xxii
 CHAPTER	
I INTRODUCTION	1
II LITERATURE REVIEW	4
Palm Oil Industry in Malaysia	4
Process of Palm Oil Extraction	4
Palm Oil By-Products as Animal Feed	
Materials	7
Nutritive Value of Palm Kernel Cake	8
Performance of Ruminants Fed PKC or PKC-Based	
Diets	19
Diary Cattle	19
Beef Cattle	20
Buffaloes	23
Goats and Sheep	24
Dietary Protein Degradation	25
Strategies for Feeding Ruminants	26



		Page
	Effects of Formaldehyde Concentrations on CP Degradability of PKC	107
	Discussion	120
V	PERFORMANCE OF BEEF CATTLE FED FORMALDEHYDE TREATED PALM KERNEL CAKE	124
	Introduction	124
	Materials and Methods	125
	Experiment 1 -Feeding Trial	126
	Experiment 2 -Digestibility Trial	128
	Results	131
	Effects of Types of PKC and Formaldehyde Treatment on Dry Matter Intake, Growth Rate and Feed Conversion Ratio in Cattle	131
	Effects of Types of PKC and Formaldehyde Treatment on Efficiency of Feed Utilization	138
	Discussion	146
VI	EFFECTS OF SUPPLEMENTING PKC-BASED DIETS WITH GRASS AND UREA ON THE PERFORMANCE AND CARCASS CHARACTERISTICS OF LAMBS	155
	Introduction	155
	Materials and Methods	157
	Animals and their Management	157
	Experimental Design	158
	Diets and Feeding Methods	159
	Measurement and Sampling Methods	159
	Slaughter and Cutting Procedures	161
	Analytical Methods	162
	Statistical Analyses	163
	Results	163
	Chemical Composition of Dietary Components	163
	Feed Intake, Growth Rate and Feed Conversion Ratio of Lambs Fed PKC Supplemented with Grass and Urea	164
	Effects of Grass and Urea Supplementation on Efficiency of Dietary Utilization	171
	Carcass Characteristics	188
	Discussion	193

	Page
VII GENERAL DISCUSSION AND CONCLUSIONS..	205
BIBLIOGRAPHY	215
APPENDIX	243
BIOGRAPHICAL VITA	245

LIST OF TABLES

Table No.		Page
1	Nutrient Compositions of Palm Kernel Cake (Based on Dry Matter)	9
2	Amino Acid Compositions of Palm Kernel Cake (g amino acid/16 g N)	11
3	Apparent Digestibility Coefficients. Digestible Energy and Metabolizable Energy of Palm Kernel Cake in the Ruminants..	13
4	Mineral Compositions of Palm Kernel Cake (Based on Dry Matter).....	17
5	Comparisons of Chemical Compositions, Gross Energy, Shell Content and Mineral Contents between PKC-Solvent Extract (PKC-S) and Expeller Pressed (PKC-E)	55
6	The Mean Values of Ruminal pH. Ammonia-N and Volatile Fatty Acids for Cattle Fed PKC-S and PKC-E	58
7	Water Intake, Fluid Rate Constant, Rumen Fluid Volume, Fluid Outflow Rate and Mean Retention Time (MRT) in Cattle Fed PKC-S or PKC-E	70
8	Dry Matter Intake (DMI), Water Intake (WI), Passage Rate Constants, Total Retention Time, Rumen Retention Time, Lower Tract Retention Time, and Transit Time of Small Particles from Rectal Sampling in Cattle Fed PKC	72
9	Degradation Parameters of Dry Matter (DM), Organic Matter (OM) and Crude Protein (CP) of PKC-S and PKC-E	77
10	Percentage of Protein Degradation of PKC-S and PKC-E Untreated and Treated with Formaldehyde at Various Treatment Times after 24 h Incubation in the Rumen of Cattle.	104

	Page
11 Percentage Protein Degradation of PKC-S and PKC-E Treated with Formaldehyde after 24 h Incubation in the Rumen of Cattle Fed PKC	105
12 Apparent Protective Effect of Formaldehyde on Protein in PKC-S and PKC-E (Percent Protection after Being Incubated in Rumen of Cattle for 24 h)	107
13 Percentage of CP Remaining in PKC-S and PKC-E after 36 h <i>In Situ</i> Incubation at Different Formaldehyde Levels	111
14 Degradation Rate and Effective Protein Degradability Estimates of Formaldehyde-Treated and Untreated PKC-S	113
15 Degradation Rate and Effective Protein Degradability Estimates of Formaldehyde-Treated and Untreated PKC-E	114
16 Relationship of Formaldehyde Treatment at Various Concentrations and By-pass Protein in PKC-S and PKC-E	116
17 Apparent Protective Effect of Formaldehyde on Protein in PKC-S and PKC-E	119
18 Effects of Formaldehyde Treatment on Protein Digestibility of PKC-S and PKC-E Estimated from <i>In Vitro</i> Pepsin Digestion (%)	120
19 Compositions of Diets	127
20 Effects of Dietary Treatment on Dry Matter Intake (DMI), Growth Rate and Feed Conversion Ratio of Cattle	134
21 Effects of PKC-S and PKC-E on Dry Matter Intake (DMI), Growth Rate and Feed Conversion Ratio of Cattle	136
22 Effects of Formaldehyde Treatment on Dry Matter Intake (DMI), Growth Rate and Feed Conversion Ratio of Cattle	137

		Page
23	Comparative Voluntary Feed Intake (VFI) and Apparent Digestibility Coefficient of Nutrients of Cattle Fed PKC-S and PKC-E Untreated and Treated with Formaldehyde	139
24	Comparative Voluntary Feed Intake (VFI) and Apparent Digestibility Coefficient of Nutrients of Cattle Fed PKC-S and PKC-E	140
25	Effects of Formaldehyde Treatment on Voluntary Feed Intake (VFI) and Apparent Digestibility Coefficient of Nutrient of Cattle	141
26	Mean Values of Gross Energy (GE), Apparent Digestibility of Gross Energy (DGE) and Total Digestible Nutrient (TDN) of Cattle Fed PKC-S and PKC-E Untreated and Treated with Formaldehyde	143
27	Mean Values of Nitrogen Retention and Nitrogen Balance of Cattle Fed PKC-S and PKC-E Untreated and Treated with Formaldehyde	145
28	Amino Acid Compositions of Rumen Microbial Protein (Bacterial and Protozoal Protein) and PKC Protein (g amino acid/16 g N)	152
29	Compositions of the Dietary Treatments	160
30	Effects of Grass Supplementation on Feed Intake, Growth Rate and Feed Conversion Ratio of Lambs Fed PKC-Based Diets	167
31	Effects of Urea Supplementation on Feed Intake, Growth Rate and Feed Conversion Ratio of Lambs Fed PKC-Based Diets	168
32	Mean Values for Initial Liveweight, Final Liveweight, Feed Intake, Growth Rate and Feed Conversion Ratio of Lambs Fed PKC Supplemented with Grass and Urea	170

	Page
33 Effects of Grass Supplementation on Intakes of Dry Matter (DM), Organic Matter (OM), Ether Extract (EE), Neutral Detergent Fibre (NDF) and Acid Detergent Fibre (ADF) and their Digestibilities in Lambs Fed PKC-Based Diets	172
34 Effects of Urea Supplementation on Intakes of Dry Matter (DM), Organic Matter (OM), Ether Extract (EE), Neutral Detergent Fibre (NDF) and Acid Detergent Fibre (ADF) and their Digestibilities in Lambs Fed PKC-Based Diets	173
35 Mean Values of Intakes of Dry Matter (DM), Organic Matter(OM), Ether Extract (EE), Neutral Detergent Fibre (NDF) and Acid Detergent Fibre (ADF) and their Digestibilities in Lambs Fed PKC Supplemented with Grass and Urea	174
36 Mean Values of Gross Energy (GE) Intake, Apparent Digestibility of GE (DGE) and Digestible Energy (DE) in Lambs Fed PKC Supplemented with Grass and Urea	177
37 Main Effects of Grass Supplementation on Nitrogen Intake (NI) and Utilization in Lambs Fed PKC-Based Diets	179
38 Main Effects of Urea Supplementation on Nitrogen Intake (NI) and Utilization in Lambs Fed PKC-Based Diets	181
39 Mean Values of Nitrogen Intake (NI) and Utilization in Lambs Fed PKC Supplemented with Grass and Urea	183
40 Effects of Grass Supplementation on Rumen pH, Concentrations and Molar Proportions of Volatile Fatty Acids (VFA) in Lambs Fed PKC-Based Diets	185
41 Effects of Urea Supplementation on Rumen pH, Concentrations and Molar Proportions of Volatile Fatty Acids (VFA) in Lambs Fed PKC-Based Diets	186

	Page
42 Mean Values for Concentrations and Molar Proportions of Volatile Fatty Acids (VFA) in Lambs Fed PKC Supplemented with Grass and Urea	187
43 Effects of Slaughter Weight, Grass Supplementation and Urea Supplementation on Dressing Percent, Ribeye Area and Backfat Thickness in Lambs Fed PKC-Based Diets	189
44 Mean Values for Dressing Percent Ribeye Area and Backfat Thickness of Lambs	191
45 Effects of Slaughter Weight, Grass Supplementation and Urea Supplementation on Wholesale Cuts of Lambs Fed PKC- Based Diets	192
46 Mean Values for Wholesale Cuts of Lambs	194

LIST OF FIGURES

Figure No.		Page
1	A Breakdown of Output Percent from Palm Oil Production ...	6
2	Ruminal pH in Cattle Fed PKC-S and PKC-E	57
3	Ruminal NH ₃ -N in Cattle Fed PKC-S and PKC-E	60
4	Total Rumen VFA Concentration of Cattle Fed PKC-S and PKC-E	61
5	Molar Percent of Acetate in the Rumen of Cattle Fed PKC-S and PKC-E	63
6	Molar Percent of Propionate in the Rumen of Cattle Fed PKC-S and PKC-E	64
7	Molar Percent of Butyrate in the Rumen of Cattle Fed PKC-S and PKC-E	65
8	Molar Percent of Other VFAs (Isobutyrate + Isovalerate + Valerate) in the Rumen of Cattle Fed PKC-S and PKC-E ...	67
9	Dilution Curve for Co in Rumen Fluid of Cattle Fed PKC-S and PKC-E	69
10	Build-up Curve for Cr from Rectal Samples of Cattle Fed PKC-S and PKC-E	71
11	Dry Matter Disappearance from PKC-S and PKC-E at Various Incubation Times in the Rumen of Cattle Fed PKC ...	73
12	Organic Matter Disappearance from PKC-S and PKC-E at Various Incubation Times in the Rumen of Cattle Fed PKC ...	74
13	Crude Protein Disappearance from PKC-S and PKC-E at Various Incubation Times in the Rumen of Cattle Fed PKC ...	75

		Page
14	Effect of Formaldehyde Levels on Protein Disappearance from PKC-S at Various Incubation Times in the Rumen of Cattle	108
15	Effect of Formaldehyde Levels on Protein Disappearance from PKC-E at Various Incubation Times in the Rumen of Cattle	109
16	By-pass Protein in PKC and Formaldehyde Levels	117
17	Dry Matter Intake of Cattle Fed PKC-S and PKC-E Untreated and Treated with Formaldehyde at Various Times of the Experimental Period	132
18	Liveweight of Cattle Fed PKC-S and PKC-E Untreated and Treated with Formaldehyde at Various Times of the Experimental Period	133
19	Dry Matter Intake of Lambs Fed PKC Supplemented with Grass and Urea at Various Times of the Experimental Period	165
20	Mean Liveweight of Lambs Fed PKC Supplemented with Grass and Urea at Various Times of the Experimental Period	166
21	Break Down of Lamb Carcass	243
22	Bone Structure of Lamb Carcass	244

LIST OF ABBREVIATIONS

ADF	- Acid detergent fibre
ADL	- Acid detergent lignin
AOAC	- Association Official Agricultural Chemists
ARC	- Agricultural Research Council
CF	- Crude fibre
CP	- Crude protein
DE	- Digestible energy
DGE	- Apparent digestibility of gross energy
DM	- Dry matter
DMI	- Dry matter intake
DMRT	- Duncan's multiple range test
DNI	- Digestible nitrogen intake
DOMI	- Digestible organic matter intake
EDTA	- Ethylenediamine tetraacetic acid
EE	- Ether extract
G	- Grass supplementation
GE	- Gross energy
HCHO	- Formaldehyde
ME	- Metabolisable energy

MRT	- Mean retention time
NDF	- Neutral detergent fibre
NFE	- Nitrogen free extract
NI	- Nitrogen intake
NRC	- National Research Council
OM	- Organic matter
PKC	- Palm kernel cake
PKC-E	- Palm kernel cake-expeller press
PKC-S	- Palm kernel cake-solvent extraction
POME	- Palm oil mill effluent
PPF	- Palm press fibre
SAS	- Statistical Analysis System
TCA	- Trichloroacetic acid
TDN	- Total digestible nutrient
TMRT	- Total mean retention time
TPKC-E	- Formaldehyde treated palm kernel cake-expeller press
TPKC-S	- Formaldehyde treated palm kernel cake-solvent extraction
TT	- Transit time
U	- Urea supplementation